
Kinematic evolution of the young local associations and the Sco-Cen complex

D. Fernández^{1,2}, F. Figueras¹ and J. Torra¹

¹ Dpt. d'Astronomia i Meteorologia, IEEC-Univ. de Barcelona, Av. Diagonal 647, E-08028 Barcelona, Spain – e-mail: david.fernandez@am.ub.es

² Observatori Astronòmic del Montsec, Consorci del Montsec, Plaça Major 1, E-25691 Àger (Lleida), Spain

Summary. In this work we propose a scenario for the history of the recent star formation (during the last 20-30 Myr) in the nearest solar neighbourhood (~ 150 pc), from the study of the spatial and kinematic properties of the members of the so-called young local associations, the Sco-Cen complex and the Local Bubble, the most important structure observed in the local interstellar medium (ISM).

1 Introduction

During the last decade, several young stellar associations have been discovered within 100 pc of Earth (see [18] for a recent review) thanks to the cross-correlation of the ROSAT and Hipparcos catalogues, combined with spectroscopy observations. The stars belonging to these young local associations (hereafter, YLA) have ages in the range from a few to several tens of Myr.

The YLA not only offer insights into the star formation process in low-density environments, but also have shed light on the substellar astrophysics, since tens of brown dwarfs have been identified in these associations. Furthermore, the study of the YLA can also yield important clues on the recent star formation in our vicinity and its consequences on the local ISM.

2 The nearest solar neighbourhood

2.1 The Local Bubble

The local interstellar medium is dominated, in the first 100 pc, by the so-called Local Bubble (LB). It was discovered at the end of the 1960s as a soft X-ray diffuse background (SXRb). Rapidly, it was recognized the existence of an anticorrelation between the X-ray background and the observed HI column density. The displacement model (see [14]) for this structure assumes that the

HI irregular local cavity is filled by an X-ray-emitting plasma, with an emission temperature of $\sim 10^6$ K and a density of $n_e \sim 0.005 \text{ cm}^{-3}$. The contours of the LB were obtained in [11] from NaI absorption measurements towards a selected set of stellar targets with Hipparcos parallaxes. These observations allowed the authors to draw maps of the neutral gas distribution in the local ISM and, in particular, to trace the contours and extension of the LB with an estimated precision of $\approx \pm 20$ pc in most directions.

Several models have been presented to explain the origin of the LB. In [4] five conceptions of the LB were reviewed. The maximum consensus is reached with a scenario where about 10-20 supernovae (SNe) formed the local cavity and, after that, 1 or 2 SNe reheated the LB a few Myr later, explaining the present temperature observed for the SXRb (see [3]).

2.2 Young local associations

A decade ago, the number of pre-main sequence stars identified near than 100 pc from the Sun was very low. Nearly all the youngest stars (< 30 Myr) studied then were located further than 140 pc, in the molecular clouds of Taurus, Chamaeleon, Lupus, Scorpius-Centaurus and R CrA, all of them regions of recent star formation. The cross-correlation of Hipparcos and ROSAT catalogues suddenly changed this view, since a few stars were identified as very young, but located closer than 100 pc, where there are not molecular clouds or star forming regions (SFR). Two scenarios were proposed to explain the existence of these young stars far away from SFR. In [15] it was defended that these stars were born in molecular clouds, but later they were ejected from them. On the other hand, in [5] it was proposed that these young stars were formed inside small molecular clouds (or *cloudlets*), which were lately dispersed in the ISM and then they are not detected at present.

During the last decade these nearby, young stars far away from molecular clouds and SFR have been identified as members of young associations, each one of them formed by a few tens of members. We have done an intensive search in the literature, and compiled all the present available data for these young associations, which are summarized in Table 1.

2.3 The Scorpius-Centaurus complex

The Scorpius-Centaurus (Sco-Cen) complex dominates the 4th galactic quadrant. It is a region of recent star formation and contains an important fraction of the most massive stars in the solar vicinity (see [17]). In the 1960s the complex was splitted into three components: Upper Scorpius (US), Upper Centaurus Lupus (UCL) and Lower Centaurus Crux (LCC). The most accepted ages for these three associations were derived in [9] from isochrones at the HR diagram: ~ 5 -6 Myr for US, ~ 14 -15 Myr for UCL and ~ 11 -12 Myr for LCC. In [2] it was suggested that this progression in age was the result of a sequence of different events of star formation which happened in the giant molecular cloud

which formed the region of Sco-Cen. These *classical* ages have been recently called in question from studies of the low mass component of the complex. The ages obtained from these studies are in the range of 8-10 Myr for US and 16-20 Myr for UCL and LCC (see [13]). The three associations show a strong velocity component in the direction away from the Sun (see Table 1), classically associated to the expansion motion of the Gould Belt (see e.g.[16]).

Table 1. Mean spatial coordinates and heliocentric velocity components of the YLA and the Sco-Cen complex. N is the number of known members of each association. Units: ξ', η', ζ' in pc; U, V, W in km s⁻¹; Age in Myr (1: from [9] and 2: from [13]).

Association	ξ'	η'	ζ'	U	V	W	Age	N
TW Hya	-21 ₍₂₂₎	-53 ₍₂₃₎	21 _(7)	-9.7 _(4.1)	-17.1 _(3.1)	-4.8 _(3.7)	~8	39
Tuc-Hor/GAYA-12	-24 ₍₂₂₎	-24 ₍₁₁₎	-34 _(8)	-10.1 _(2.4)	-20.7 _(2.3)	-2.5 _(3.8)	20-30	52
β Pic-Cap	-9 ₍₂₇₎	-5 ₍₁₄₎	-15 ₍₁₀₎	-10.8 _(3.4)	-15.9 _(1.2)	-9.8 _(2.5)	~12	33
ϵ Cha	-47 _(8)	-80 ₍₁₄₎	-25 _(5)	-8.6 _(3.6)	-18.6 _(0.8)	-9.3 _(1.7)	5-15	16
η Cha	-33 _(2)	-80 _(5)	-34 _(2)	-12.2 _(0.0)	-18.1 _(0.9)	-10.1 _(0.5)	<10	18
HD 141569	-77 _(3)	10 _(8)	64 _(8)	-5.4 _(1.5)	-15.6 _(2.6)	-4.4 _(0.8)	2-8	5
Ext. R CrA	-97 ₍₄₄₎	-1 _(4)	-30 ₍₁₆₎	-0.1 _(6.4)	-14.8 _(1.4)	-10.1 _(3.3)	10-15	59
AB Dor	6 ₍₂₁₎	4 ₍₁₇₎	-12 ₍₁₆₎	-7.4 _(3.2)	-27.4 _(3.2)	-12.9 _(6.4)	75-150	40
US	-134	-20	52	-6.7 _(5.9)	-16.0 _(3.5)	-8.0 _(2.7)	5-6 ¹ 8-10 ²	120
UCL	-119	-67	31	-6.8 _(4.6)	-19.3 _(4.7)	-5.7 _(2.5)	14-15 ¹ 16-20 ²	221
LCC	-62	-100	10	-8.2 _(5.1)	-18.6 _(7.3)	-6.4 _(2.6)	11-12 ¹	180

3 Kinematic evolution of the young stellar component

In this section we present the results of our kinematic study of the YLA and the Sco-Cen complex. This study is based on the integration back in time of the orbits of these associations, allowing us to study their origin and their possible influence on the nearest ISM during the last million years. The integration of the equations of motion has been done using a realistic galactic potential which includes an axisymmetric component, the spiral arm perturbation and the central bar contribution (see [1] and [7]).

3.1 Orbits back in time: the origin of the YLA

Studying the orbits back in time for the YLA, we have found as the most gaudy trend a spatial concentration in the 1st galactic quadrant (see Fig. 1), in such a way that those stars belonging to the YLA were concentrated in a region of about 60 x 100 x 40 pc at their birth instant, whereas at present they are spread inside a region of 120 x 130 x 140 pc in size (except for AB

Dor, which is clearly older than the rest of YLA; see [7]). Therefore, we can affirm that a star formation process was triggered in this region of the 1st galactic quadrant, between 5 and 15 Myr ago, producing the YLA. What was the mechanism which triggered this star formation process? One possibility is the explosion of one or several close SNe, which should belong to the most important concentration of O and B stars in our vicinity: the Sco-Cen complex.

It is then of great interest to determine the instant when the minimum distance between LCC (the closest Sco-Cen association to the YLA during the last few Myr) and the YLA occurred. For 3 YLA (η Cha, ϵ Cha and Ext. R CrA), minimum distances of ~ 16 , 23 and 73 pc are obtained for $t \sim -9$ Myr. No clear minima in distance to LCC are found for the other YLA, but distances of only a few tens of pc are obtained.

3.2 The YLA and the Local Bubble

We have found that the trajectories back in time for the YLA have nearly crossed the center of the Local Bubble (LB) during the last ~ 5 Myr (see Fig. 4 in [11]). Studying the stellar content of the YLA (see [7] for details), we conclude that it is possible that one or a few of these associations have sheltered a SN in the recent past (the last 10 Myr). On the other hand, there is direct evidence for an explosion of a SN at a distance of ~ 30 pc from the Sun, ~ 5 Myr ago (see [10]). Taking into account the trajectories back in time of the Sco-Cen associations that we have computed, it is not probable that this SN could exploded in one of them, since not even LCC has approached as much to the solar neighbourhood in the past. Then, several pieces of the same puzzle seem to support the existence of a SN explosion in the nearest solar neighbourhood (~ 30 pc) ~ 5 Myr ago, from a parent star belonging to one of the YLA, probably Tuc-Hor/GAYA or Ext. R CrA. This close and recent SN would have been responsible for the reheating of the gas inside the LB needed to achieve the observed temperatures in the soft X-ray diffuse background at present.

4 A scenario for the recent star formation in our vicinity

To complete the scenario outlined in the previous section, we should study the origin of the Sco-Cen associations. Several scenarios have been proposed to explain the origin of this region (see [13]), but a careful analysis (see [7]) allows to conclude that the most promising is the impact with a spiral arm. This scenario is supported by several theoretical and observational facts (see [7] and [8]). Taking this as an hypothesis, the history of the star formation in the nearest solar neighbourhood during the last few tens of million years would have been written as follows. 30 Myr ago, the giant molecular cloud parent of Sco-Cen was placed in a position of the galactic plane with coordinates $(X', Y') \sim (-400, 1200)$ pc. The arriving of the potential minimum of the




Fig. 1. Orbits back in time on the galactic plane to $t = -30$ Myr for the YLA and the Sco-Cen associations, in the LSR reference frame (RF; top) and in the RF which is rotating at the same velocity of the spiral arms (bottom). ξ' and X' point to the galactic center, and η' , Y' point to the direction of the galactic rotation. The thick-dashed line shows the position of the minimum of the spiral potential (adopted from [6]). In thin-dashed line, the position of the phase of the spiral structure $\psi = 10^\circ$.

inner spiral arm triggered the star formation in the region, disturbing at the same time the cloud motion, which began to move with a velocity vector directed to the galactic antirotation and away from the galactic center (see Fig. 1; just the expected motion after an interaction with the spiral arm for a position outside the corotation radius). The compression due to the spiral arm did not necessarily trigger the star formation in the whole cloud, but perhaps only in those regions with the largest densities. This would be favoured by the smaller relative velocity between the shock wave and the RSR (12-13 km s⁻¹, see [7]). The regions where the star formation began must be those which generate UCL, LCC and, probably, the Tuc-Hor/GAYA association, which

were born nearly at the same time, about 16-20 Myr ago. The stellar winds of the first massive stars began to compress the gas of the neighbouring regions, maybe producing their fragmentation in small molecular clouds which moved away from the central region of the parent molecular cloud. 9 Myr ago, a SN in LCC could trigger the star formation in these small molecular clouds, giving birth the majority of the YLA, as we have seen in the previous section. The stellar winds of the just born stars rapidly expelled the remaining gas from these small clouds (the *cloudlets* proposed by [5]), completely erasing every trace of them and promoting that we do not observe gas in these regions at present time. Later, as it was proposed in [12], the shock front of a SN in UCL would have triggered the star formation in US about 6 Myr ago. Only 1.5 Myr ago, the most massive star in US would have exploded as SN and its shock front would be reaching now the molecular cloud of ρ Oph, triggering there the beginning of the star formation process. This would complete our scenario for the recent star formation in the nearest solar neighbourhood.

Acknowledgements. This work has been supported by the CICYT under contracts AYA2003-07736 and AYA2006-15623-C02-02.

References

1. R. Asiain, F. Figueras, J. Torra: A&A **350**, 434 (1999)
2. A. Blaauw: ARA&A **2**, 213 (1964)
3. D. Breitschwerdt, D.P. Cox: Is the Local Bubble Dead?. In: *How does the Galaxy work? A Galactic Tertulia with Don Cox and Ron Reynolds*, ed by E.J. Alfaro, E. Pérez, J. Franco (A&SS Library, vol. 315, 2004), 391–401
4. D.P. Cox: Modeling the Local Bubble. In: *Proceedings of the IAU Colloquium 166: The Local Bubble and Beyond*, ed by D. Breitschwerdt, M.J. Freyberg, J. Trümper (Lecture Notes in Physics, vol. 506, 1998), 121–126
5. E.D. Feigelson: ApJ **468**, 306 (1996)
6. D. Fernández, F. Figueras, J. Torra: A&A **372**, 833 (2001)
7. D. Fernández, F. Figueras, J. Torra: A&A, submitted (2007)
8. D. Fernández: Estructura espacial y cinemática de la componente estelar joven en el entorno solar. PhD Thesis, Universitat de Barcelona, Barcelona (2005)
9. E. de Geus, P.T. de Zeeuw, J. Lub: A&A **216**, 44 (1989)
10. K. Knie, G. Korschinek, T. Faestermann et al.: Phys. Rev. Lett. **81**, 18 (1999)
11. R. Lallement, B.Y. Welsh, J.L. Vergely, F. Crifo, D.M. Sfeir: A&A **411**, 447 (2003)
12. T. Preibisch, H. Zinnecker: AJ **117**, 2381 (1999)
13. M.J. Sartori, J.R.D. Lépine, W.S. Dias: A&A **404**, 913 (2003)
14. S.L. Snowden, D.P. Cox, D. McCammon, W.T. Sanders: ApJ **354**, 211 (1990)
15. M. Sterzik, R. Durisen: A&A **304**, L9 (1995)
16. J. Torra, D. Fernández, F. Figueras: A&A **359**, 82 (2000)
17. P.T. de Zeeuw, R. Hoogerwerf, J.H.J. Bruijne et al.: AJ **117**, 354 (1999)
18. B. Zuckerman, I. Song: ARA&A **42**, 685 (2004)